B.Sc. Semester-II Core Course-III (CC-III) Organic Chemistry-I



IV. Aromatic Hydrocarbons

2. Benzene and Criteria for Aromaticity



Dr. Rajeev Ranjan University Department of Chemistry Dr. Shyama Prasad Mukherjee University, Ranchi

IV Aromatic Hydrocarbons

10 Lectures

Aromaticity: Hückel's rule, aromatic/anti-aromatic/non-aromatic character of arenes, cyclic carbocations/carbanions and heterocyclic compounds with suitable examples.

Electrophilic aromatic substitution: Halogenation, Nitration, Sulphonation and Friedel-Craft's alkylation/acylation with their mechanism. Directing effects of mono-functional groups.

Coverage:

- 1. Benzene and Criteria for Aromaticity
- 2. Hückel's rule, aromatic/anti-aromatic/non-aromatic character of arenes

Benzene (C₆H₆) is not "Cyclohexatriene"!



Each sp² hybridized C in the ring has an unhybridized *p*-orbital perpendicular to the ring which overlaps around the ring

Criteria for Aromaticity

- 1. Structure must be cyclic
- 2. Unhybridized p orbital in continuous cyclic system
- 3. Able to adopt a planar geometry
- 4. Fulfills Huckel's Rule and includes (4n + 2) electrons n = 0, 1, 2, 3,

4.....

(the 2, 6, 10, 14....electrons)



 Initially, all cyclic conjugated hydrocarbons were proposed to be aromatic

•However, cyclobutadiene is so reactive that it dimerizes before it can be isolated

•Cyclooctatetraene adds Br₂ readily.

•Look at Molecular Orbitals (MOs) to explain aromaticity in benzenelike molecules

Catalytic hydrogenation of benzene and various cyclohexenes



Constructing π -Molecular Orbitals diagram

• π molecular orbitals are the sideways overlap of *p* orbitals

•*p* orbitals have 2 lobes. Plus (+) and minus (-) indicate the opposite phases of the wave function, <u>not</u> electrical charge

•When lobes overlap constructively, (+ and +, or - and -) a bonding MO is formed

•When + and - lobes overlap, waves cancel out and a node forms; antibonding MO



π -Molecular Orbitals for Benzene

- •Six overlapping *p*-orbitals must form six molecular orbitals
- •Three will be three bonding, three antibonding orbitals
- •Lowest energy MO will have all bonding interactions, no nodes
- •As energy of MO increases, the number of nodes increases
- •System symmetric so 2 pairs of degenerate orbitals



π - Molecular Orbital Energy Diagram for Benzene

6 atomic orbitals - 6 molecular orbitals

System symmetric so 2 pairs of degenerate orbitals



The six electrons fill three bonding pi orbitals.

All bonding orbitals are filled ("closed shell"), an extremely stable arrangement (Aromatic Stabilization).

π - Molecular Orbital Energy Diagram for Cyclobutadiene



If cyclobutadiene adopted a planar geometry - two of the molecular orbitals would each have a single unpaired electron - very unstable. Applies to any (4 n) system Cyclobutadiene is ANTIAROMATIC





Dr. Rajeev Ranjan

raene- Non-aromatic due to 1055 of plananty

Thank You



Dr. Rajeev Ranjan University Department of Chemistry Dr. Shyama Prasad Mukherjee University, Ranchi